

Requirements for compatibility between local and multipartite quantum states

Sergey Bravyi*

*Institute for Quantum Information,
California Institute of Technology,
Pasadena, CA, 91125, USA*

February 1, 2008

Abstract

We consider a partial trace transformation which maps a multipartite quantum state to collection of local density matrices. We call this collection a *mean field state*. For the Hilbert spaces $(\mathbb{C}^2)^{\otimes n}$ and $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^4$ the necessary and sufficient conditions under which a mean field state is compatible with at least one multipartite pure state are found. Compatibility of mean field states with more general classes of multipartite quantum states is discussed.

1 Introduction and main results

Consider a system consisting of n quantum particles and denote $\mathcal{H}^{(i)}$ a Hilbert space of states of the i -th particle. A state of the whole system can be most exhaustively described by a vector $|\Psi\rangle$ from the Hilbert space

$$\mathcal{H} = \mathcal{H}^{(1)} \otimes \dots \otimes \mathcal{H}^{(n)}$$

(a pure state). However, sometimes we need not such a detailed description. For example if we are interested only in expectation values of one-particle observables, it is sufficient to keep track of just n reduced density matrices $\rho^{(1)} = \text{tr}_{i \neq 1}(|\Psi\rangle\langle\Psi|), \dots, \rho^{(n)} = \text{tr}_{i \neq n}(|\Psi\rangle\langle\Psi|)$ describing the states of the individual particles (here the subscript i shows the particles to be traced out). This is some kind of mean field description because we completely ignore all types of correlations (both quantum and classical) between the different particles.

An arbitrary collection of n density matrices $(\rho^{(1)}, \dots, \rho^{(n)})$ will be referred to as a *mean field state*. Naturally, some mean field states can not be represented as partial traces of any pure state. For example let us take $n = 2$ and $\mathcal{H}^{(1)} = \mathcal{H}^{(2)}$. From the Schmidt decomposition we learn that the density matrices $\rho^{(1)}$ and $\rho^{(2)}$ arising from a pure state must share the same set of the eigenvalues. If this is not the case, we can say that a mean field state $(\rho^{(1)}, \rho^{(2)})$ is not compatible with a pure multipartite state. Conversely, if the eigenvalues of $\rho^{(1)}$ and $\rho^{(2)}$ match, we can always find a pure state $|\Psi\rangle$ such that $\rho^{(1)} = \text{tr}_2(|\Psi\rangle\langle\Psi|)$ and $\rho^{(2)} = \text{tr}_1(|\Psi\rangle\langle\Psi|)$.

*e-mail: serg@cs.caltech.edu

The main purpose of this paper is to find the necessary and sufficient conditions for compatibility between mean field states and multipartite pure states in the case of three or more particles where Schmidt decomposition can not be applied. For any Hilbert space $\mathcal{S}$ denote $D(\mathcal{S})$ a set of all Hermitian, non-negative operators with unital trace (a set of density matrices). Now we can formulate our first problem.

Problem 1. *Given a multipartite system $\mathcal{H} = \mathcal{H}^{(1)} \otimes \dots \otimes \mathcal{H}^{(n)}$, find a set of all mean field states $(\rho^{(1)}, \dots, \rho^{(n)})$, $\rho^{(1)} \in D(\mathcal{H}^{(1)}), \dots, \rho^{(n)} \in D(\mathcal{H}^{(n)})$ which are compatible with at least one multipartite pure state $|\Psi\rangle \in \mathcal{H}$.*

Let us denote the set to be found M_{pure} . Its formal definition is

$$M_{pure} = \{(\rho^{(1)}, \dots, \rho^{(n)}) : \rho^{(i)} = \text{tr}_{j \neq i}(|\Psi\rangle\langle\Psi|), |\Psi\rangle \in \mathcal{H}, \langle\Psi|\Psi\rangle = 1\}.$$

The compatibility problem may be generalized to the case when a state of the whole system is a mixed one with a specified set of the eigenvalues. Denote $\text{eig}(\rho)$ a string of eigenvalues of an operator ρ (to avoid ambiguity, let us agree that the eigenvalues are arranged in the decreasing order). Denote also $D(\mathcal{S}, \lambda) = \{\rho \in D(\mathcal{S}) : \text{eig}(\rho) = \lambda\}$ a set of density matrices with a specified spectrum λ . Our second problem is following.

Problem 2. *Given a multipartite system $\mathcal{H} = \mathcal{H}^{(1)} \otimes \dots \otimes \mathcal{H}^{(n)}$ and a string of eigenvalues λ , find a set of all mean field states $(\rho^{(1)}, \dots, \rho^{(n)})$, $\rho^{(1)} \in D(\mathcal{H}^{(1)}), \dots, \rho^{(n)} \in D(\mathcal{H}^{(n)})$ which are compatible with at least one multipartite mixed state $\rho \in D(\mathcal{H}, \lambda)$.*

Let us denote the set to be found $M(\lambda)$. Its formal definition is

$$M(\lambda) = \{(\rho^{(1)}, \dots, \rho^{(n)}) : \rho^{(i)} = \text{tr}_{j \neq i}(\rho), \rho \in D(\mathcal{H}, \lambda)\}.$$

The Problem 2 can be reduced to the Problem 1 by introducing one additional particle (ancilla) with a space of states $\mathcal{H}$. One suffices to find the set M_{pure} for the extended system and then find the subset of M_{pure} corresponding to the fixed eigenvalues of the ancilla. In that sense the two problems are equivalent.

The solution of the Problem 1 for an arbitrary n is known only for the simplest case $\mathcal{H}^{(i)} = \mathbb{C}^2$ for all i , see [6]. This is a system of n spins $\frac{1}{2}$ or, equivalently, of n qubits. Let $(\rho^{(1)}, \dots, \rho^{(n)})$ be an arbitrary mean field state of the system. Denote $\lambda^{(i)}$ the lowest eigenvalue of i -th qubit's density matrix $\rho^{(i)}$. By definition, $\lambda^{(i)} \in [0, \frac{1}{2}]$. Then $(\rho^{(1)}, \dots, \rho^{(n)}) \in M_{pure}$ if and only if for each i in the range $1, 2, \dots, n$ one has

$$\lambda^{(i)} \leq \sum_{j \neq i} \lambda^{(j)}. \quad (1)$$

For instance, if $n = 2$, one gets: $\lambda^{(1)} \leq \lambda^{(2)}$, $\lambda^{(2)} \leq \lambda^{(1)}$, that is $\lambda^{(1)} = \lambda^{(2)}$ as it should be. For $n = 3$ one gets the triangle inequalities: $\lambda^{(1)} \leq \lambda^{(2)} + \lambda^{(3)}$, $\lambda^{(2)} \leq \lambda^{(3)} + \lambda^{(1)}$, $\lambda^{(3)} \leq \lambda^{(1)} + \lambda^{(2)}$. For real spin systems, such as magnetics, we have $n \gg 1$ and the inequalities (1) impose practically no restrictions on the mean field states. However these inequalities may be important in quantum information theory, in particular in the study of multipartite analogs of the Schmidt decomposition [1, 2] and polynomial invariants under local unitary transformations [3, 4, 5]. In Section 2 we describe an alternative proof of (1) obtained independently from the work [6].

We have solved the Problem 2 only for a bipartite system and only in the simplest case $\mathcal{H}^{(1)} = \mathcal{H}^{(2)} = \mathbb{C}^2$. This is a system of two qubits. Let us label the qubits by letters A and B.

Consider an arbitrary mean field state (ρ^A, ρ^B) . Denote λ^A and λ^B the lowest eigenvalues of the density matrices ρ^A and ρ^B . By definition, $\lambda^A, \lambda^B \in [0, \frac{1}{2}]$. Let $\lambda = \{\lambda_1, \lambda_2, \lambda_3, \lambda_4\}$ be an arbitrary string of real non-negative numbers, such that $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \lambda_4$ and $\sum_{i=1}^4 \lambda_i = 1$. In Section 3 we will show that $(\rho^A, \rho^B) \in M(\lambda)$ if and only if

$$\begin{aligned} \lambda^A &\geq \lambda_4 + \lambda_3, \\ \lambda^B &\geq \lambda_4 + \lambda_3, \\ \lambda^A + \lambda^B &\geq 2\lambda_4 + \lambda_3 + \lambda_2, \\ |\lambda^A - \lambda^B| &\leq \min\{\lambda_1 - \lambda_3, \lambda_2 - \lambda_4\}. \end{aligned} \tag{2}$$

This result also provides a solution of the Problem 1 for $n = 3$ with $\mathcal{H}^{(1)} = \mathcal{H}^{(2)} = \mathbb{C}^2$ and $\mathcal{H}^{(3)} = \mathbb{C}^4$.

Although we can not attack the Problem 2 in the general case, we can solve its simplified version, namely find the convex envelope [8] of the set $M(\lambda)$:

$$\text{Conv}(M(\lambda)) = \{(\rho^{(1)}, \dots, \rho^{(n)}) : \rho^{(i)} = \text{tr}_{j \neq i}(\rho), \rho \in \text{Conv}(D(\mathcal{H}, \lambda))\},$$

where $\text{Conv}(D(\mathcal{H}, \lambda))$ is the convex envelope of the set $D(\mathcal{H}, \lambda)$. By definition, any density matrix $\rho \in \text{Conv}(D(\mathcal{H}, \lambda))$ can be represented as $\rho = \sum_{\alpha} p_{\alpha} U_{\alpha} \rho_0 U_{\alpha}^{\dagger}$ for some probabilities $p_{\alpha} \geq 0$, $\sum_{\alpha} p_{\alpha} = 1$, unitary operators U_{α} and some density matrix $\rho_0 \in D(\mathcal{H}, \lambda)$. The set $\text{Conv}(D(\mathcal{H}, \lambda))$ is important for example for description of interaction between the system and the *classical* environment.

The simplified Problem 2 is considered in Section 4. We show that it can be reduced to the analogous *classical* problem concerning multipartite probability distributions. To describe this reduction, let us consider as an example a bipartite system of Alice and Bob, $\mathcal{H} = \mathcal{H}^A \otimes \mathcal{H}^B$. Let λ_j^A be the j -th element of the string $\text{eig}(\rho^A)$ and λ_j^B be the j -th element of the string $\text{eig}(\rho^B)$ (one can order the eigenvalues by an arbitrary way; this ambiguity will not affect the answer). Let d^A and d^B be the dimensions of the Hilbert spaces $\mathcal{H}^A$ and $\mathcal{H}^B$. For any bipartite probability distribution $p^{AB}(i, j)$ of integer variables $i \in [1, d^A]$ and $j \in [1, d^B]$ denote p^{AB} a string of $d^A \cdot d^B$ probabilities $p^{AB}(1, 1), \dots, p^{AB}(1, d^B), \dots, p^{AB}(d^A, 1), \dots, p^{AB}(d^A, d^B)$ ordered by an arbitrary way (this ambiguity will not affect the answer). In Section 4 we will show that

$$\text{Conv}(M(\lambda)) = \{(\rho^A, \rho^B) : \lambda_i^A = \sum_{j=1}^{d^B} p^{AB}(i, j), \lambda_j^B = \sum_{i=1}^{d^A} p^{AB}(i, j), p^{AB} \prec \lambda\}, \tag{3}$$

where $p^{AB} \prec \lambda$ means that the string p^{AB} is majorized by the string λ (the majorization relation is briefly discussed in Section 4). A set of probability distributions $p^{AB}(i, j)$ satisfying $p^{AB} \prec \lambda$ is known to be a convex one, and a probability distribution $p^{AB}(i, j)$ is an extremal point if and only if $p^{AB} = \lambda$ up to some permutation of the elements. This set of probability distributions is thus a classical analogue of $\text{Conv}(D(\mathcal{H}, \lambda))$. The generalization of equality (3) to the case of three or more parties is straightforward.

2 Mean field states for arrays of qubits.

In this section we consider a multipartite system consisting of qubits (spins $\frac{1}{2}$) and prove the equality (1). Let $(\rho^{(1)}, \dots, \rho^{(n)})$ be an arbitrary mean field state of n qubit system. By

appropriate unitary local operators we can always bring the matrices $\rho^{(i)}$ to the standard form

$$\rho^{(i)} = \lambda^{(i)}|1\rangle\langle 1| + (1 - \lambda^{(i)})|0\rangle\langle 0|, \quad i \in [1, n], \quad (4)$$

where $\lambda^{(i)}$ is the *lowest* eigenvalue of $\rho^{(i)}$, that is $\lambda^{(i)} \in [0, \frac{1}{2}]$. The proof of (1) consists of two parts.

(a) Suppose that $(\rho^{(1)}, \dots, \rho^{(n)}) \in M_{\text{pure}}$, i.e. $\rho^{(i)} = \text{tr}_{j \neq i}(|\Psi\rangle\langle\Psi|)$ for some pure state $|\Psi\rangle \in (\mathbb{C}^2)^{\otimes n}$. Let us prove, for example, the inequality $\lambda^{(n)} \leq \sum_{j=1}^{n-1} \lambda^{(j)}$. If n -qubit system is described by the pure state $|\Psi\rangle$, the qubits $1, \dots, n-1$ alone are described by the density matrix

$$\rho = \text{tr}_n(|\Psi\rangle\langle\Psi|). \quad (5)$$

Taking into account (4) we can write

$$\sum_{j=1}^{n-1} \lambda^{(j)} = \langle\Psi| \sum_{j=1}^{n-1} \Pi_1[j] |\Psi\rangle = \text{tr}(\rho(\sum_{j=1}^{n-1} \Pi_1[j])), \quad (6)$$

where $\Pi_1 = |1\rangle\langle 1|$ is a one-qubit projector and the designation $\Pi_1[j]$ means that the operator Π_1 is applied to the j -th qubit. Note that the operator $\sum_{j=1}^{n-1} \Pi_1[j]$ is diagonal in the standard qubit basis and counts a number of 1's in a basis vector. Its only zero eigenvalue corresponds to the state $|00 \dots 0\rangle$. For that reason we will introduce an auxiliary operator

$$H = \sum_{j=1}^{n-1} \Pi_1[j] + |00 \dots 0\rangle\langle 00 \dots 0| \quad (7)$$

Clearly, all eigenvalues of H are greater or equal to unity, i.e. $H \geq I$. As the whole n -qubit system is described by the pure state $|\Psi\rangle$, the positive eigenvalues of the density matrices ρ and $\rho^{(n)}$ coincide. In particular the highest eigenvalue of ρ is equal to the highest eigenvalue of $\rho^{(n)}$ which is $1 - \lambda^{(n)}$. It means that $\langle\psi|\rho|\psi\rangle \leq (1 - \lambda^{(n)})\langle\psi|\psi\rangle$ for any $|\psi\rangle \in (\mathbb{C}^2)^{\otimes n-1}$. Thus we conclude that

$$\sum_{j=1}^{n-1} \lambda^{(j)} = \text{tr}(\rho H) - \langle 00 \dots 0|\rho|00 \dots 0\rangle \geq \text{tr}(\rho) - (1 - \lambda^{(n)}) = \lambda^{(n)}, \quad (8)$$

which is the desired inequality.

(b) We should prove that any mean field state satisfying the inequalities (1) does arise from some n -qubit pure state. Obviously one and the same mean field state can arise from many pure states, so we have some freedom in choosing them. We will consider only *even pure states* (EPS) which are the states of the following form:

$$|\Psi\rangle = \sum_{x \in B_{\text{even}}^n} c_x |x\rangle, \quad x = (x_1 \dots x_n), \quad x_i \in \{0, 1\}. \quad (9)$$

Here B_{even}^n is a set of all binary strings with the even number of 1's. The EPS's span the linear subspace of dimension 2^{n-1} .

For any choice of the amplitudes c_x in (9) the reduced density matrices $\rho^{(i)}$ are diagonal in the $\{|0\rangle, |1\rangle\}$ basis and look as

$$\rho^{(i)} = \left(\sum_{x \in B_{\text{even}}^n : x_i=0} |c_x|^2 \right) |0\rangle\langle 0| + \left(\sum_{x \in B_{\text{even}}^n : x_i=1} |c_x|^2 \right) |1\rangle\langle 1|, \quad i \in [1, n]. \quad (10)$$

Thus to guarantee that the matrices $\rho^{(i)}$ have the standard form (4) we should only require that

$$\langle 0|\rho^{(i)}|0\rangle \geq \langle 1|\rho^{(i)}|1\rangle, \quad i \in [1, n]. \quad (11)$$

Besides, the set of all mean field states which arise from the EPS's is a convex one. Indeed, if we take an arbitrary pair of mean field states $(\rho_0^{(1)}, \dots, \rho_0^{(n)})$ and $(\rho_1^{(1)}, \dots, \rho_1^{(n)})$ such that

$$\rho_0^{(i)} = \text{tr}_{j \neq i}(|\Psi_0\rangle\langle\Psi_0|) \text{ and } \rho_1^{(i)} = \text{tr}_{j \neq i}(|\Psi_1\rangle\langle\Psi_1|), \quad i \in [1, n],$$

for some EPS's

$$|\Psi_0\rangle = \sum_{x \in B_{even}^n} c_{x,0} |x\rangle \text{ and } |\Psi_1\rangle = \sum_{x \in B_{even}^n} c_{x,1} |x\rangle,$$

then for any $t \in [0, 1]$ we can realize the linear combination $(1-t)(\rho_0^{(1)}, \dots, \rho_0^{(n)}) + t(\rho_1^{(1)}, \dots, \rho_1^{(n)})$ by the EPS

$$|\Psi_t\rangle = \sum_{x \in B_{even}^n} \sqrt{(1-t)|c_{x,0}|^2 + t|c_{x,1}|^2} |x\rangle.$$

Consequently a set of all mean filed states which arise from EPS's and satisfy the additional requirements (11) is also a convex one.

The mean field state written in the standard form (4) can be identified with a real vector $\vec{\lambda} \in \mathbb{R}^n$ defined as $\vec{\lambda} = (\lambda^{(1)}, \dots, \lambda^{(n)})$. The previous statement can be rephrased by saying that the vectors $\vec{\lambda}$ arising from EPS's constitute some convex subset in $\mathbb{R}^n$ which we will denote Λ_{EPS} . By definition, $0 \leq \lambda^{(i)} \leq \frac{1}{2}$, so that Λ_{EPS} is a convex subset of the n -dimensional cube I with the edge length $1/2$. Denote V the set of all 2^n vertices of I , i.e. the vectors $\vec{v} = (\epsilon_1, \dots, \epsilon_n)$, $\epsilon \in \{0, 1/2\}$. There are exactly n vertices $\vec{v} \in V$ which violate the inequalities (1). These are the vertices containing exactly one $1/2$ component. All other $2^n - n$ vertices satisfy (1) and contain either zero or greater than one $1/2$ components. It will be convenient to introduce a special designation to distinguish the vertices of the last type:

$$V^* = \{\vec{v} \in V : \vec{v} = (\epsilon_1 \dots \epsilon_n), \quad \epsilon_i \leq \sum_{j \neq i}^n \epsilon_j, \quad \epsilon_i \in \{0, \frac{1}{2}\}, \quad i \in [1, n]\}. \quad (12)$$

We claim that any vector from V^* can be realized by an EPS, i.e. $V^* \subset \Lambda_{EPS}$. Indeed, the vectors

$$(0 \dots 0), \left(\frac{1}{2} \frac{1}{2} 0 \dots 0\right), \left(\frac{1}{2} \frac{1}{2} \frac{1}{2} 0 \dots 0\right), \dots \left(\frac{1}{2} \dots \frac{1}{2}\right) \in V^*$$

can be realized by the EPS's as follows:

$$\begin{aligned} (0 \dots 0) & : |\Psi\rangle = |0 \dots 0\rangle, \\ \left(\frac{1}{2} \frac{1}{2} 0 \dots 0\right) & : |\Psi\rangle = 2^{-\frac{1}{2}}(|00\rangle + |11\rangle) \otimes |0 \dots 0\rangle, \\ \left(\frac{1}{2} \frac{1}{2} \frac{1}{2} 0 \dots 0\right) & : |\Psi\rangle = 2^{-1}(|000\rangle + |011\rangle + |101\rangle + |110\rangle) \otimes |0 \dots 0\rangle, \\ & \dots \\ \left(\frac{1}{2} \dots \frac{1}{2}\right) & : |\Psi\rangle = 2^{-\frac{n-1}{2}} \sum_{x \in B_{even}^n} |x\rangle. \end{aligned} \quad (13)$$

All other vectors from V^* can be realized by EPS's applying appropriate qubit permutations to the states listed above. We know that Λ_{EPS} is a convex set, so that any vector from the convex

envelope $Conv(V^*)$ also can be realized by an EPS, i.e. $Conv(V^*) \subset \Lambda_{EPS}$. To conclude the proof we will need a simple geometrical lemma which claims that the set $Conv(V^*)$ coincides with those part of the cube I which is specified by the inequalities (1):

$$Conv(V^*) = \{\vec{\lambda} \in I : \lambda^{(i)} \leq \sum_{j \neq i} \lambda^{(j)}, \quad i \in [1, n]\}. \quad (14)$$

We encourage the reader to verify this statement for $n = 2, 3$ by drawing a picture. The proof of (14) for an arbitrary n is given in the Appendix A. Thus for any $\vec{\lambda} \in I$ such that $\lambda^{(i)} \leq \sum_{j \neq i} \lambda^{(j)}$, $i \in [1, n]$ we have proven that $\vec{\lambda} \in Conv(V^*) \subset \Lambda_{EPS}$, so that $\vec{\lambda}$ can be realized by the EPS. It means that any mean field state satisfying (1) can be realized by some n -qubit pure state.

Note that (14) implies also a converse inclusion $\Lambda_{EPS} \subset Conv(V^*)$ because we already know that inequalities (1) are necessary for compatibility of the mean field state with a pure state. It means that $\Lambda_{EPS} = Conv(V^*)$.

3 Mean field states for two qubits

In this section we consider a bipartite system composed of two qubits and prove that the inequalities (2) are necessary for compatibility of a mean field state (ρ^A, ρ^B) with at least one mixed state from $D(\mathbb{C}^4, \lambda)$. A proof that (2) is also a sufficient condition for compatibility is placed in Appendix B.

The combinations of λ_i 's standing in the righthand sides of the inequalities (2) can be easily found if we consider two *classical* bits and joint probability distributions of two bits with the specified set of probabilities $\{\lambda_1, \lambda_2, \lambda_3, \lambda_4\}$ which can be arbitrarily assigned to the events 00, 01, 10, and 11 (this is a classical analog of the set $D(\mathbb{C}^4, \lambda)$). Generally there are $4!$ such distributions. In this case the reduced density matrices ρ^A, ρ^B become the partial one-bit probability distributions and λ^A, λ^B become the minimal values of ρ^A, ρ^B . In these settings the inequalities (2) are the best estimates on the quantities $\lambda^A, \lambda^B, \lambda^A + \lambda^B$, and $|\lambda^A - \lambda^B|$ as can be explicitly verified. The nontrivial result is that this estimates also hold in the quantum case and the set $M(\lambda)$ is completely specified by them.

As we will see below, the strict proof of the inequality $|\lambda^A - \lambda^B| \leq \min\{\lambda_1 - \lambda_3, \lambda_2 - \lambda_4\}$ is rather formidable. However in one particular case it can be proved without any calculations. Suppose that the minimum here is zero, i.e. $\lambda_1 = \lambda_3$ or $\lambda_2 = \lambda_4$. Then the spectrum λ involves an eigenvalue which is degenerated with multiplicity three or four. In this case we can represent any density matrix $\rho \in D(\mathbb{C}^4, \lambda)$ as a linear combination of the identity operator and a projector on some pure state. Obviously the term proportional to identity operator just shifts the spectrums of ρ^A and ρ^B by the same constant. So it doesn't contribute to the combination $\lambda^A - \lambda^B$. The term proportional to a projector on a pure state produces the reduced density matrices ρ^A and ρ^B with the same spectrum according to the Schmidt theorem. So we conclude that $\lambda^A = \lambda^B$.

Suppose that $(\rho^A, \rho^B) \in M(\lambda)$, i.e. $\rho^A = tr_B(\rho)$, $\rho^B = tr_A(\rho)$ for some $\rho \in D(\mathbb{C}^4, \lambda)$. We should prove that λ^A, λ^B , and λ satisfy four inequalities (2). Let us represent the matrices ρ^A, ρ^B as

$$\rho^A = \lambda^A \Pi^A + (1 - \lambda^A)(I - \Pi^A), \quad \rho^B = \lambda^B \Pi^B + (1 - \lambda^B)(I - \Pi^B), \quad (15)$$

where Π^A , Π^B are the projectors on the eigenstates of ρ^A , ρ^B with the lowest eigenvalue, so that $\lambda^A = \text{tr}(\rho\Pi^A[A])$, $\lambda^B = \text{tr}(\rho\Pi^B[B])$. As usual the qubit acted on by one-qubit operator is indicated in the square brackets.

Suppose now that O is an arbitrary two-qubit hermitian operator with the eigenvalues $O_1 \geq O_2 \geq O_3 \geq O_4$. We claim that

$$\inf_{\eta \in D(\mathbb{C}^4, \lambda)} \text{tr}(O\eta) = \lambda_1 O_4 + \lambda_2 O_3 + \lambda_3 O_2 + \lambda_4 O_1. \quad (16)$$

To verify this statement one can consider the variation $\delta\eta = i[h, \eta]$ where $h^\dagger = h$ is an arbitrary infinitesimal hermitian operator. This variation is obviously consistent with the constraint $\eta \in D(\mathbb{C}^4, \lambda)$. Suppose that the minimum in (16) is achieved at the matrix $\eta^* \in D(\mathbb{C}^4, \lambda)$. Then we have

$$\delta \text{tr}(O\eta^*) = i \text{tr}(O[h, \eta^*]) = i \text{tr}(h[\eta^*, O]) = 0 \quad (17)$$

for any h . It means that $[\eta^*, O] = 0$ so that η^* and O are diagonal in the same basis. Therefore there is a permutation $\sigma \in S_4$ such that

$$\text{tr}(O\eta^*) = \lambda_1 O_{\sigma(4)} + \lambda_2 O_{\sigma(3)} + \lambda_3 O_{\sigma(2)} + \lambda_4 O_{\sigma(1)}. \quad (18)$$

It is easy to verify that the optimal permutation is always $\sigma(i) = i$ and thus the equality (16) is proven.

Let us substitute $O = \Pi^A[A]$ into the equality (16). In this case we have $\{O_1, O_2, O_3, O_4\} = \{1, 1, 0, 0\}$ and thus

$$\lambda^A = \text{tr}(\rho\Pi^A[A]) \geq \inf_{\eta \in D(\mathbb{C}^4, \lambda)} \text{tr}(\eta\Pi^A[A]) = \lambda_4 + \lambda_3. \quad (19)$$

By the same way substituting $O = \Pi^B[B]$ and $O = \Pi^A[A] + \Pi^B[B]$ into (16) we get the inequalities $\lambda^B \geq \lambda_4 + \lambda_3$ and $\lambda^A + \lambda^B \geq 2\lambda_4 + \lambda_2 + \lambda_3$.

Unfortunately the last inequality of (2), that is

$$|\lambda^A - \lambda^B| \leq \min\{\lambda_1 - \lambda_3, \lambda_2 - \lambda_4\}, \quad (20)$$

can not be proved using the same idea (the analogous method allows to get the estimate $|\lambda^A - \lambda^B| \leq \lambda_1 - \lambda_4$ which is too weak). The proof of (20) presented below is a bit formidable.

Let us define polarization vectors for the qubits A and B:

$$a^\alpha[\eta] = \text{tr}(\sigma^\alpha[A]\eta), \quad b^\alpha[\eta] = \text{tr}(\sigma^\alpha[B]\eta), \quad (21)$$

where $\sigma^x, \sigma^y, \sigma^z$ are the Pauli matrices and $\eta \in D(\mathbb{C}^4)$ is an arbitrary density matrix. Denote also $a = \sqrt{a^\alpha a^\alpha}$ and $b = \sqrt{b^\alpha b^\alpha}$ the absolute magnitudes of the polarization vectors. We can express the eigenvalues λ^A and λ^B in terms of polarizations as $\lambda^A = \frac{1}{2}(1 - a[\rho])$ and $\lambda^B = \frac{1}{2}(1 - b[\rho])$. Introduce a functional

$$F[\eta] = b[\eta] - a[\eta], \quad \eta \in D(\mathbb{C}^4), \quad (22)$$

such that $|\lambda^A - \lambda^B| = \frac{1}{2}|F[\rho]|$. The inequality (20) is equivalent to the following one:

$$\sup_{\eta \in D(\mathbb{C}^4, \lambda)} F[\eta] \leq 2 \min\{\lambda_1 - \lambda_3, \lambda_2 - \lambda_4\} \quad (23)$$

Note that an application of the qubit swap transformation changes the sign of F and doesn't change the eigenvalues of η so the supremum (23) can not be negative. Let $\eta^* \in D(\mathbb{C}^4, \lambda)$ be some density matrix at which this supremum is achieved (may be it is not unique). Consider the variation $\delta\eta = i[h, \eta^*]$ where $h^\dagger = h$ is an arbitrary infinitesimal hermitian operator. This variation is consistent with the constraint $\eta \in D(\mathbb{C}^4, \lambda)$. We must require that $\delta F[\eta^*] = 0$.

Suppose first that both polarization vectors $a^\alpha \equiv a^\alpha[\eta^*]$ and $b^\alpha \equiv b^\alpha[\eta^*]$ are non-zero. Denote $\hat{a}^\alpha = a^\alpha/a$ and $\hat{b}^\alpha = b^\alpha/b$ the normalized polarization vectors. The equation $\delta F[\eta^*] = 0$ then can be written as

$$\text{tr}(h [\eta^*, \hat{a}^\alpha \sigma^\alpha[A] - \hat{b}^\alpha \sigma^\alpha[B]]) = 0. \quad (24)$$

As h may be an arbitrary hermitian operator we conclude that

$$\hat{a}^\alpha [\sigma^\alpha[A], \eta^*] = \hat{b}^\alpha [\sigma^\alpha[B], \eta^*]. \quad (25)$$

Let us parameterize the density matrix η^* as follows:

$$\eta^* = \frac{1}{4}(I + a^\alpha \sigma^\alpha[A] + b^\alpha \sigma^\alpha[B] + A_{\alpha,\beta} \sigma^\alpha[A] \sigma^\beta[B]), \quad (26)$$

where $A_{\alpha,\beta}$ is some 3x3 real matrix. Using the commutators $[\sigma^\alpha, \sigma^\beta] = 2i\epsilon^{\alpha\beta\gamma} \sigma^\gamma$ we can transform the equation (25) into

$$\hat{a}^\gamma A_{\delta,\alpha} \epsilon^{\gamma\delta\beta} = \hat{b}^\gamma A_{\beta,\delta} \epsilon^{\gamma\delta\alpha}. \quad (27)$$

As the functional $F[\eta]$ is invariant under one-qubit unitary transformations, we can assume that both qubits are polarized along z -axis, i.e. $\hat{a} = \hat{b} = (0, 0, 1)$. Under this assumption one can easily check that the most common solution of the equation (27) looks as

$$A = \begin{bmatrix} s & t & 0 \\ t & -s & 0 \\ 0 & 0 & r \end{bmatrix} \quad (28)$$

for some real numbers s, t, r . Substituting it into (26) we can represent the matrix η^* in the basis $\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$ of $\mathbb{C}^4$ as follows

$$\eta^* = \frac{1}{4} \begin{bmatrix} 1+a+b+r & 0 & 0 & \bar{z} \\ 0 & 1+a-b-r & 0 & 0 \\ 0 & 0 & 1-a+b-r & 0 \\ z & 0 & 0 & 1-a-b+r \end{bmatrix}, \quad (29)$$

where $z = 2(s + it)$. Note that η^* is a block diagonal matrix so its eigenvalues $\{\lambda_1^*, \lambda_2^*, \lambda_3^*, \lambda_4^*\}$ can be easily found:

$$\begin{aligned} \lambda_1^* &= \frac{1}{4}(1+a-b-r), & \lambda_3^* &= \frac{1}{4}(1+r+\sqrt{(a+b)^2+|z|^2}), \\ \lambda_2^* &= \frac{1}{4}(1-a+b-r), & \lambda_4^* &= \frac{1}{4}(1+r-\sqrt{(a+b)^2+|z|^2}). \end{aligned} \quad (30)$$

Recall that $\eta^* \in D(\mathbb{C}^4, \lambda)$ so that the eigenvalues λ_i^* coincide up to some permutation with the eigenvalues λ_i . Although this permutation somehow depends upon a, b, r, z (recall that the

eigenvalues λ_i are ordered in decreasing way) we need not to consider all possible cases. Instead we can notice that

$$\max_{\sigma \in S^4} \min\{|\lambda_{\sigma(1)} - \lambda_{\sigma(3)}|, |\lambda_{\sigma(2)} - \lambda_{\sigma(4)}|\} = \min\{\lambda_1 - \lambda_3, \lambda_2 - \lambda_4\}, \quad (31)$$

where the maximum is taken over all permutations of four elements. Note also that the lefthand side is invariant under the substitution $\lambda_i \rightarrow \lambda_i^*$. Thus we conclude that

$$\min\{|\lambda_1^* - \lambda_2^*|, |\lambda_3^* - \lambda_4^*|\} \leq \min\{\lambda_1 - \lambda_3, \lambda_2 - \lambda_4\}. \quad (32)$$

Taking λ_i^* from the table (30) we arrive to the inequality

$$\frac{1}{2} \min\{|a - b|, \sqrt{(a + b)^2 + |z|^2}\} \leq \min\{\lambda_1 - \lambda_3, \lambda_2 - \lambda_4\}. \quad (33)$$

The minimum in the lefthand side is equal to $|a - b|$ regardless of the parameter z . Consequently

$$b - a \leq 2 \min\{\lambda_1 - \lambda_3, \lambda_2 - \lambda_4\}, \quad (34)$$

which proves the inequality (23).

Suppose now that at the supremum of $F[\eta]$ one of the polarization vectors becomes zero. The only non-trivial situation is $a^\alpha[\eta^*] = 0$. In this case the requirement $\delta F[\eta^*] = 0$ under the variation $\delta\eta = i[h, \eta^*]$ implies that $\delta a^\alpha[\eta^*] = 0$ for all three components α . By analogy with (25) this requirement implies that

$$[\sigma^\alpha[A], \eta^*] = 0, \quad \alpha = x, y, z. \quad (35)$$

It means that η^* commutes with any operator acting only on the qubit A. This is possible only if η^* has the form $\eta^* = (I/2) \otimes \eta^B$ for some one-qubit density matrix η^B . Thus the eigenvalues of η^* are

$$\lambda_1 = \lambda_2 = \frac{1}{4}(1 + b), \quad \lambda_3 = \lambda_4 = \frac{1}{4}(1 - b). \quad (36)$$

Note that $\eta^* \in D(\mathbb{C}^4, \lambda)$ only if $\lambda_1 = \lambda_2$ and $\lambda_3 = \lambda_4$ so the considered situation is not a generic one. Nevertheless we have $F[\eta^*] = b = 2(\lambda_1 - \lambda_3) = 2(\lambda_2 - \lambda_4)$ so that the estimate (23) is valid in this case also.

4 Connection with multipartite probability distributions

In this section we consider a bipartite system with local Hilbert spaces $\mathcal{H}^A$ and $\mathcal{H}^B$ of arbitrary dimensions. We show that a set

$$\text{Conv}(M(\lambda)) = \{(\rho^A, \rho^B) : \rho^A = \text{tr}_B(\rho), \rho^B = \text{tr}_A(\rho), \rho \in \text{Conv}(D(\mathcal{H}, \lambda))\} \quad (37)$$

can be described in terms of multipartite probability distributions according to (3). The restriction that the system is bipartite is not essential and is needed only to simplify the designations. Using the arguments presented below, one can easily formulate and prove a multipartite analogue of (3). We start by collecting all relevant facts concerning the majorization relation.

Consider arbitrary strings of non-negative real numbers $p = (p_1, \dots, p_d)$ and $q = (q_1, \dots, q_d)$ such that $\sum_{i=1}^d p_i = \sum_{i=1}^d q_i = 1$. Let us arrange the elements of p and q in decreasing order.

Denote $p_i^\downarrow$ the i -th element of p from above and $q_i^\downarrow$ the i -th element of q from above, such that $p_1^\downarrow \geq p_2^\downarrow \geq \dots \geq p_d^\downarrow$ and $q_1^\downarrow \geq q_2^\downarrow \geq \dots \geq q_d^\downarrow$. The string p is said to be majorized by the string q , if for each k in the range $1, \dots, d-1$ we have

$$\sum_{i=1}^k p_i^\downarrow \leq \sum_{i=1}^k q_i^\downarrow. \quad (38)$$

This relation is denoted as $p \prec q$. We will need three well known facts given below:

(i) $p \prec q$ if and only if $p_i = \sum_{\sigma \in S_d} t_\sigma q_{\sigma(i)}$, $i = 1, \dots, d$, where S_d is the symmetric group, $\sigma \in S_d$ is a permutation of d elements, $t_\sigma \geq 0$ and $\sum_{\sigma \in S_d} t_\sigma = 1$. The existence of such decomposition is guaranteed by Birkhoff and von Neumann decomposition of a doubly stochastic matrix into a convex combination of permutation matrices.

(ii) Let $\mathcal{S}$ be an arbitrary Hilbert space. Consider any pair of density matrices $\rho, \eta \in D(\mathcal{S})$. Then $\text{eig}(\rho) \prec \text{eig}(\eta)$ if and only if $\rho = \sum_\alpha t_\alpha U_\alpha \eta U_\alpha^\dagger$, where U_α are some unitary operators on $\mathcal{S}$, $t_\alpha \geq 0$ and $\sum_\alpha t_\alpha = 1$, see [7].

(iii) Let $\mathcal{S}$ be an arbitrary Hilbert space. For any $\rho \in D(\mathcal{S})$ and for any orthonormal basis in $\mathcal{S}$, the string of diagonal elements of ρ is majorized by the eigenvalues of ρ , i.e. $\text{diag}(\rho) \prec \text{eig}(\rho)$.

Note that according to the property (ii), the requirement $\rho \in \text{Conv}(D(\mathcal{H}, \lambda))$ is equivalent to $\rho \in D(\mathcal{H})$ and $\text{eig}(\rho) \prec \lambda$, so we can rewrite (37) as

$$\text{Conv}(M(\lambda)) = \{(\rho^A, \rho^B) : \rho^A = \text{tr}_B(\rho), \rho^B = \text{tr}_A(\rho), \rho \in D(\mathcal{H}), \text{eig}(\rho) \prec \lambda\}. \quad (39)$$

Now we are ready to prove the result (3). The proof consists of two parts.

(a) Let (ρ^A, ρ^B) be an arbitrary mean field state such that

$$\lambda_i^A = \sum_{j=1}^{d^B} p^{AB}(i, j), \quad \lambda_j^B = \sum_{i=1}^{d^A} p^{AB}(i, j), \quad (40)$$

for some bipartite probability distribution $p^{AB}(i, j)$ satisfying $p^{AB} \prec \lambda$. Consider the eigenvectors decomposition of the density matrices ρ^A and ρ^B :

$$\rho^A = \sum_{i=1}^{d^A} \lambda_i^A |i_A\rangle \langle i_A|, \quad \rho^B = \sum_{j=1}^{d^B} \lambda_j^B |j_B\rangle \langle j_B|, \quad (41)$$

where $|i_A\rangle \in \mathcal{H}^A$, $|j_B\rangle \in \mathcal{H}^B$. Define a bipartite density matrix $\rho \in D(\mathcal{H})$ according to

$$\rho = \sum_{i=1}^{d^A} \sum_{j=1}^{d^B} p^{AB}(i, j) |i_A\rangle \langle i_A| \otimes |j_B\rangle \langle j_B|. \quad (42)$$

By definition, $\rho^A = \text{tr}_B(\rho)$, $\rho^B = \text{tr}_A(\rho)$, and $\text{eig}(\rho) = p^{AB} \prec \lambda$, so that $(\rho^A, \rho^B) \in \text{Conv}(M(\lambda))$, see (39).

(b) Now suppose that $(\rho^A, \rho^B) \in \text{Conv}(M(\lambda))$, i.e. that $\rho^A = \text{tr}_B(\rho)$, $\rho^B = \text{tr}_A(\rho)$ for some $\rho \in D(\mathcal{H})$ such that $\text{eig}(\rho) \prec \lambda$, see (39). Consider the eigenvectors decomposition (41) and define a bipartite probability distribution $p^{AB}(i, j)$ according to

$$p^{AB}(i, j) = \langle i_A, j_B | \rho | i_A, j_B \rangle, \quad i \in [1, d^A], \quad j \in [1, d^B]. \quad (43)$$

Clearly, this probability distribution satisfies the requirements (40). By definition, in the basis $\{|i_A, j_B\rangle\}_{i,j}$ we have $p^{AB} = \text{diag}(\rho)$, so the property (iii) tells us that $p^{AB} \prec \text{eig}(\rho)$. Summarizing, we have: $p^{AB} \prec \text{eig}(\rho) \prec \lambda$. Transitivity of majorization relation implies that $p^{AB} \prec \lambda$. This completes the proof of the equality (3).

For an arbitrary string of eigenvalues λ , let us denote $P(\lambda)$ the set of all bipartite probability distributions $p^{AB}(i, j)$ such that $p^{AB} = \lambda$ up to some permutation. The set $P(\lambda)$ is a classical analogue of the set $D(\mathcal{H}, \lambda)$. The property (i) tells us that $p^{AB} \prec \lambda$ if and only if $p^{AB} \in \text{Conv}(P(\lambda))$. Thus the reduction of the Problem 3 to the analogous classical problem corresponds to the substitutions $(\rho^A, \rho^B) \rightarrow (\lambda_i^A, \lambda_j^B)$, $\rho \rightarrow p^{AB}(i, j)$, $D(\mathcal{H}, \lambda) \rightarrow P(\lambda)$. The partial traces corresponds to taking the average over one variable.

The inclusion $(\rho^{(1)}, \dots, \rho^{(n)}) \in \text{Conv}(M(\lambda))$ may be regarded as a necessary condition for inclusion $(\rho^{(1)}, \dots, \rho^{(n)}) \in M(\lambda)$. It provides us some knowledge about the sets $M(\lambda)$ and M_{pure} for an arbitrary multipartite system. For example consider a system of three particles A, B, and C. Let (ρ^A, ρ^B, ρ^C) be an arbitrary mean field state. The requirements

$$\begin{aligned} (\rho^A, \rho^B) &\in \text{Conv}(M(\text{eig}(\rho^C))), \\ (\rho^B, \rho^C) &\in \text{Conv}(M(\text{eig}(\rho^A))), \\ (\rho^C, \rho^A) &\in \text{Conv}(M(\text{eig}(\rho^B))) \end{aligned} \tag{44}$$

are necessary for inclusion $(\rho^A, \rho^B, \rho^C) \in M_{\text{pure}}$. If A, B, and C are the qubits, one can check that these requirements are also sufficient. Meanwhile if $\mathcal{H}^A = \mathcal{H}^B = \mathbb{C}^2$ and $\mathcal{H}^C = \mathbb{C}^4$, one can build an explicit example which shows that they are not sufficient. Indeed, take $\text{eig}(\rho^A) = \{0.6, 0.4\}$, $\text{eig}(\rho^B) = \{0.5, 0.5\}$, and $\text{eig}(\rho^C) = \{0.3, 0.3, 0.3, 0.1\}$. One can verify that $\text{eig}(\rho^A \otimes \rho^B) \prec \text{eig}(\rho^C)$, $\text{eig}(\rho^B \otimes \rho^C) \prec \text{eig}(\rho^A)$, and $\text{eig}(\rho^C \otimes \rho^A) \prec \text{eig}(\rho^B)$, so that (44) are satisfied. However $(\rho^A, \rho^B, \rho^C) \notin M_{\text{pure}}$ because there is another necessary requirement which is violated by this mean field state, see Section 3.

5 Conclusions

We found the necessary and sufficient conditions under which a collection of density matrices (a mean field state) can be produced by the partial trace operations from a multipartite pure state for the system of n qubits and for the system $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^4$. These conditions may be described by linear inequalities on the eigenvalues of local density matrices. We also found the necessary and sufficient conditions under which a mean field state can be produced by the partial trace operations from a multipartite mixed state whose eigenvalues are majorized by a given real vector.

6 Acknowledgements

I would like to thank David DiVincenzo and Alexei Kitaev for advices on majorization technique. I am also very grateful to Alexei Kitaev for thoroughly reading the manuscript and for many suggestions which helped to improve the paper. This work was supported by NWO-Russia collaboration program.

7 Appendix A

The purpose of this section is to prove the equality (14). Denote $\hat{I}$ the subset of the cube I which is specified by the inequalities (1), i.e.

$$\hat{I} = \{\vec{\lambda} \in \mathbb{R}^n : \lambda^{(i)} \leq \sum_{j \neq i} \lambda^{(j)}, \quad 0 \leq \lambda^{(i)} \leq \frac{1}{2}, \quad i \in [1, n]\}. \quad (45)$$

Clearly I is a convex set. The extremal point of $\hat{I}$ is a vector $\vec{\lambda} \in \hat{I}$ for which some $m \geq n$ of the inequalities (45) specifying $\hat{I}$ become the equalities and the corresponding system of m equations has a unique solution (thus having a rank n). Let us prove that if $\vec{\lambda} \in \hat{I}$ is an extremal point of $\hat{I}$, then $\vec{\lambda} \in V^*$. Because any convex set is a convex envelope of its extremal points, it will imply that $\hat{I} \subset \text{Conv}(V^*)$.

To each extremal point $\vec{\lambda} \in \hat{I}$ we can assign a pair of integers (p, q) where $p \in [0, n]$ is the number of equalities $\lambda^{(i)} = \sum_{j \neq i} \lambda^{(j)}$ which are satisfied at $\vec{\lambda}$ and $q \in [0, n]$ is the number of equalities $\lambda^{(i)} = 0$, $\lambda^{(i)} = \frac{1}{2}$ which are satisfied at $\vec{\lambda}$. By definition $p + q \geq n$ but we can assume that $p + q = n$ because at each extremal point there are only n independent equations. Now we are ready to list all extremal points of $\hat{I}$ and check that they all belong to V^* .

(a) The extremal points of the type $(p, n - p)$, $p \geq 3$

We have the equalities $\lambda^{(i)} = \sum_{j \neq i} \lambda^{(j)}$, $\lambda^{(k)} = \sum_{j \neq k} \lambda^{(j)}$, $\lambda^{(l)} = \sum_{j \neq l} \lambda^{(j)}$ for some $i \neq k \neq l$.

They are compatible with the inequalities $\lambda_j \geq 0$, $j \in [1, n]$ only if $\vec{\lambda} = (0 \dots 0)$.

(b) The extremal points of the type $(2, n - 2)$.

The equalities $\lambda^{(i)} = \sum_{j \neq i} \lambda^{(j)}$, $\lambda^{(k)} = \sum_{j \neq k} \lambda^{(j)}$ and the inequalities $\lambda_j \geq 0$, $j \in [1, n]$ are compatible only if $\lambda^{(i)} = \lambda^{(k)}$ and $\lambda^{(j)} = 0$ for $j \neq i$ or $j \neq k$. Thus we have either $\lambda^{(i)} = \lambda^{(k)} = 0$ or $\lambda^{(i)} = \lambda^{(k)} = \frac{1}{2}$. The extremal points of both types belongs to V^* .

(c) The extremal points of the type $(1, n - 1)$.

Suppose that $\lambda^{(i)} = \sum_{j \neq i} \lambda^{(j)}$ for some $i \in [1, n]$. If we choose the equality $\lambda^{(i)} = 0$ then $\lambda^{(j)} = 0$ for all $j \neq i$. If we choose the equality $\lambda^{(i)} = \frac{1}{2}$ then we must choose $\lambda^{(k)} = \frac{1}{2}$ for some $k \neq i$ and $\lambda^{(j)} = 0$ for $j \neq i$ and $j \neq k$. If we do not choose neither $\lambda^{(i)} = 0$ nor $\lambda^{(i)} = \frac{1}{2}$ then for each $j \neq i$ we must choose either $\lambda^{(j)} = 0$ or $\lambda^{(j)} = \frac{1}{2}$. This choice is compatible with the equality $\lambda^{(i)} = \sum_{j \neq i} \lambda^{(j)}$ only if $\lambda^{(i)} = 0$ or $\lambda^{(i)} = \frac{1}{2}$. In all these cases we arrive to the extremal points which have been already listed in (a) or (b).

(d) The extremal points of the type $(0, n)$.

The extremal point of this type is a vertex of the cube I which satisfies the inequalities (1). According to definition (12) any point of this type belongs to V^* .

So all extremal points of $\hat{I}$ belong to V^* and thus $\hat{I} \subset \text{Conv}(V^*)$. The inclusion $\text{Conv}(V^*) \subset \hat{I}$ is obvious because $V^* \subset \hat{I}$ and $\hat{I}$ is a convex set. Thus the equality $\hat{I} = \text{Conv}(V^*)$ is proven.

8 Appendix B

In this section we show that if a mean field state of two qubits (ρ^A, ρ^B) satisfies the inequalities (2) for some eigenvalue string $\lambda = (\lambda_1, \lambda_2, \lambda_3, \lambda_4)$, then it can be represented as $\rho^A = \text{tr}_B(\rho)$, $\rho^B = \text{tr}_A(\rho)$ for some two qubit density matrix $\rho \in D(\mathbb{C}^4, \lambda)$, i.e. that $(\rho^A, \rho^B) \in M(\lambda)$. It is convenient to draw a region Ω defined by the inequalities (2) on the plane (λ^A, λ^B) for a fixed string λ , see FIG. 1. We will describe the desired matrix $\rho \in D(\mathbb{C}^4, \lambda)$ explicitly for each point




Figure 1: The region Ω on the plane (λ^A, λ^B) specified by the inequalities (2) for $\lambda = (0.7, 0.2, 0.1, 0)$. For an arbitrary λ the coordinates of the vertices O, A, B, C, D are: $O = (\frac{1}{2}, \frac{1}{2})$, $A = (\frac{1}{2}, \frac{1}{2} - \min[\lambda_1 - \lambda_3, \lambda_2 - \lambda_4])$, $B = (\min[\lambda_1 + \lambda_4, \lambda_2 + \lambda_3], \lambda_3 + \lambda_4)$, $C = (\lambda_2 + \lambda_4, \lambda_3 + \lambda_4)$, $D = \frac{1}{2}(2\lambda_4 + \lambda_3 + \lambda_2, 2\lambda_4 + \lambda_3 + \lambda_2)$.

inside Ω . It is sufficient to consider only the case $\lambda^A \geq \lambda^B$ because the constraint $\rho \in D(\mathbb{C}^4, \lambda)$ is invariant under the permutation of the qubits. Also we will assume that $\lambda_1 - \lambda_3 \geq \lambda_2 - \lambda_4$. The opposite case may be investigated similarly. We will describe three independent families of the density matrices $\rho \in D(\mathbb{C}^4, \lambda)$ corresponding to three triangles OCD , OAB , and OBC on FIG. 1. Let us define two qubit states

$$\begin{aligned}
|\Psi_1\rangle &= 2^{-\frac{1}{2}} [\sqrt{1-\alpha} |0,0\rangle + \sqrt{1+\alpha} |1,1\rangle], \\
|\Psi_2\rangle &= 2^{-\frac{1}{2}} [\sqrt{1+\alpha} |0,0\rangle - \sqrt{1-\alpha} |1,1\rangle], \\
|\Psi_3\rangle &= 2^{-\frac{1}{2}} [\sqrt{1-\beta} |0,1\rangle + \sqrt{1+\beta} |1,0\rangle], \\
|\Psi_4\rangle &= 2^{-\frac{1}{2}} [\sqrt{1+\beta} |0,1\rangle - \sqrt{1-\beta} |1,0\rangle].
\end{aligned} \tag{46}$$

For any parameters $\alpha, \beta \in [-1, 1]$ this is an orthonormal basis in $\mathbb{C}^4$, i.e. $\langle \Psi_i | \Psi_j \rangle = \delta_{i,j}$. Consider the density matrix

$$\rho = \lambda_1 |\Psi_1\rangle \langle \Psi_1| + \lambda_2 |\Psi_4\rangle \langle \Psi_4| + \lambda_3 |\Psi_3\rangle \langle \Psi_3| + \lambda_4 |\Psi_2\rangle \langle \Psi_2| \in D(\mathbb{C}^4, \lambda). \tag{47}$$

The reduced density matrices $\rho^A = tr_B(\rho)$ and $\rho^B = tr_A(\rho)$ have the lowest eigenvalues

$$\lambda^A = \frac{1}{2} \{1 - |\alpha(\lambda_1 - \lambda_4) - \beta(\lambda_2 - \lambda_3)|\}, \quad \lambda^B = \frac{1}{2} \{1 - |\alpha(\lambda_1 - \lambda_4) + \beta(\lambda_2 - \lambda_3)|\}. \tag{48}$$

One can easily verify that the region of parameters $0 \leq \alpha \leq 1$, $0 \leq \beta \leq \alpha$ is mapped to the triangle OCD on FIG. 1. Next consider the density matrix

$$\rho = \lambda_1 |\Psi_1\rangle \langle \Psi_1| + \lambda_2 |\Psi_4\rangle \langle \Psi_4| + \lambda_3 |\Psi_2\rangle \langle \Psi_2| + \lambda_4 |\Psi_3\rangle \langle \Psi_3| \in D(\mathbb{C}^4, \lambda). \tag{49}$$

The reduced density matrices $\rho^A = \text{tr}_B(\rho)$ and $\rho^B = \text{tr}_A(\rho)$ have the lowest eigenvalues

$$\lambda^A = \frac{1}{2} \{1 - |\alpha(\lambda_1 - \lambda_3) - \beta(\lambda_2 - \lambda_4)|\}, \quad \lambda^B = \frac{1}{2} \{1 - |\alpha(\lambda_1 - \lambda_3) + \beta(\lambda_2 - \lambda_4)|\}. \quad (50)$$

One can easily verify that the region of parameters $0 \leq \beta \leq 1$, $\frac{\lambda_2 - \lambda_4}{\lambda_1 - \lambda_3} \beta \leq \alpha \leq \beta$ is mapped to the triangle OAB on FIG. 1. Now let us define two qubit states

$$\begin{aligned} |\tilde{\Psi}_1\rangle &= 2^{-\frac{1}{2}} [\sqrt{1-\alpha} |0,0\rangle + \sqrt{1+\alpha} |1,1\rangle], \\ |\tilde{\Psi}_2\rangle &= 2^{-\frac{1}{2}} [\sqrt{1+\alpha} |\phi_0,0\rangle - \sqrt{1-\alpha} |M_1,1\rangle], \\ |\tilde{\Psi}_3\rangle &= 2^{-\frac{1}{2}} [\sqrt{1-\alpha} |\phi_0,1\rangle + \sqrt{1+\alpha} |\phi_1,0\rangle], \\ |\tilde{\Psi}_4\rangle &= 2^{-\frac{1}{2}} [\sqrt{1+\alpha} |0,1\rangle - \sqrt{1-\alpha} |1,0\rangle], \end{aligned} \quad (51)$$

where $|\phi_0\rangle = \cos(\frac{\varphi}{2})|0\rangle + \sin(\frac{\varphi}{2})|1\rangle$ and $|\phi_1\rangle = -\sin(\frac{\varphi}{2})|0\rangle + \cos(\frac{\varphi}{2})|1\rangle$. For any parameters $\alpha \in [-1, 1]$ and $\varphi \in [0, \pi]$ this is an orthonormal basis in $\mathbb{C}^4$, i.e. $\langle \tilde{\Psi}_i | \tilde{\Psi}_j \rangle = \delta_{i,j}$. Consider the density matrix

$$\rho = \lambda_1 |\tilde{\Psi}_1\rangle \langle \tilde{\Psi}_1| + \lambda_2 |\tilde{\Psi}_4\rangle \langle \tilde{\Psi}_4| + \lambda_3 |\tilde{\Psi}_3\rangle \langle \tilde{\Psi}_3| + \lambda_4 |\tilde{\Psi}_2\rangle \langle \tilde{\Psi}_2| \in D(\mathbb{C}^4, \lambda). \quad (52)$$

The reduced density matrices $\rho^A = \text{tr}_B(\rho)$ and $\rho^B = \text{tr}_A(\rho)$ have the lowest eigenvalues

$$\lambda^A = \frac{1}{2} \left(1 - |\alpha| \cdot \sqrt{\lambda_{12}^2 + \lambda_{34}^2 + 2\lambda_{12}\lambda_{34}\cos\varphi} \right), \quad \lambda^B = \frac{1}{2} [1 - |\alpha| \cdot (\lambda_1 + \lambda_2 - \lambda_3 - \lambda_4)], \quad (53)$$

where $\lambda_{12} = \lambda_1 - \lambda_2$ and $\lambda_{34} = \lambda_3 - \lambda_4$. One can easily verify that the region of parameters $0 \leq \alpha \leq 1$, $0 \leq \varphi \leq \pi$ is mapped to the triangle OBC on FIG. 1. Our goal is achieved.

References

- [1] H. A. Carteret, A. Higuchi, and A. Sudbery, “Multipartite generalization of the Schmidt decomposition“, LANL preprint archive quant-ph/0006125.
- [2] A. Acin, A. Andrianov, L. Costa, E. Jane, J.I. Latorre, and R. Tarrach, “Generalized Schmidt decomposition and classification of three-quantum-bit states“, LANL preprint archive quant-ph/0003050.
- [3] K.G.H. Vollbrecht and R.F.Werner, “Entanglement measures under symmetry“, LANL preprint archive quant-ph/0010095
- [4] A.Sudbery, “On local invariants of pure three-qubit states“, *J.Phys.A* **34**, 643-652 (2001), also available at LANL preprint archive quant-ph/0001116
- [5] H.Barnum and N.Linden, “Monotones and invariants for multi-particle quantum states“, LANL preprint archive quant-ph/0103155
- [6] A. Higuchi, A. Sudbery, and J. Szulc, “One-qubit reduced states of a pure many-qubit state: polygon inequalities“, LANL preprint archive quant-ph/0209085

- [7] A. Wehrl, “How chaotic is a state of a quantum system?”, *Reports on Mathematical Physics*, vol. 6, pp. 15–28, 1974.
- [8] Recall that for any subset X of a linear space $\mathcal{L}$, the convex envelope $\text{Conv}(X) \subset \mathcal{L}$ is a set of all linear combinations $\sum_{\alpha} p_{\alpha} x_{\alpha}$, where $x_{\alpha} \in X$, p_{α} are non-negative real numbers, and $\sum_{\alpha} p_{\alpha} = 1$. We will often say about convex sets of mean field states assuming that their linear combinations are defined as $t(\rho_1^{(1)}, \dots, \rho_1^{(n)}) + (1 - t)(\rho_0^{(1)}, \dots, \rho_0^{(n)}) \equiv (t\rho_1^{(1)} + (1 - t)\rho_0^{(1)}, \dots, t\rho_1^{(n)} + (1 - t)\rho_0^{(n)})$.